

Isometries of infinite dimensional Hilbert geometries

Bas Lemmens

*School of Mathematics, Statistics & Actuarial Science
University of Kent, Canterbury, Kent CT2 7NX, UK
B.Lemmens@kent.ac.uk*

Mark Roelands* and Marten Wortel†

*Unit for BMI, North-West University
Private Bag X6001-209, Potchefstroom 2520, South Africa
*mark.roelands@gmail.com
†marten.wortel@gmail.com*

Received 30 September 2015

Accepted 22 February 2017

Published 7 April 2017

In this paper we extend two classical results concerning the isometries of strictly convex Hilbert geometries, and the characterisation of the isometry groups of Hilbert geometries on finite dimensional simplices, to infinite dimensions. The proofs rely on a mix of geometric and functional analytic methods.

Keywords: Hilbert geometries; isometries; projective linear homomorphisms.

AMS Subject Classification: Primary: 58B20; Secondary: 22F50, 46B04

1. Introduction

In [10] Hilbert introduced a collection of metric spaces that are natural deformations of finite dimensional real hyperbolic spaces. Although Hilbert limited his construction to finite dimensions, it has a straightforward extension to infinite dimensional spaces. Indeed, let Ω be a convex subset of a (not necessarily finite-dimensional) real vector space Y , and suppose that for each $x \neq y \in \Omega$ the straight line ℓ_{xy} through x and y has the property that $\Omega \cap \ell_{xy}$ is an open and bounded line segment in ℓ_{xy} . In that case one can define Hilbert's metric on Ω as follows. For $x \neq y$ in Ω , let x' and y' be the end-points of the segment $\ell_{xy} \cap \Omega$ such that x is between x' and y , and y is between y' and x . Now *Hilbert's metric on Ω* is given by

$$\delta_H(x, y) := \log[x', x, y, y'],$$

where

$$[x', x, y, y'] := \frac{|x' - y| |y' - x|}{|x' - x| |y' - y|}$$

is the *cross-ratio*, and $\delta_H(x, x) = 0$ for all $x \in \Omega$. The metric space (Ω, δ_H) is usually called the *Hilbert geometry on Ω* . In particular, the open unit ball in an infinite dimensional Hilbert space equipped with $\frac{1}{2}\delta_H$, is precisely Klein's model of the infinite dimensional hyperbolic space. A recent extensive overview of the theory of Hilbert geometries can be found in [21].

The isometries between finite dimensional Hilbert geometries are well understood. They have been studied intensively in the past decade by Bosché [3], de la Harpe [7], Lemmens and Walsh [15], Matveev and Troyanov [17], Speer [22], and Walsh [24]. The purpose of this paper is to analyse the isometries of infinite dimensional geometries on strictly convex domains and infinite dimensional simplices. To date there are only a few works on infinite dimensional Hilbert geometries. We should mention the work [18] by Molnár in which the group of Hilbert's metric isometries on the projective domain of the cone of positive self-adjoint operators on a complex Hilbert space is determined, and [4, 19] in which the isometries of infinite dimensional hyperbolic space are studied.

It is well known that Hilbert's metric has important applications in the analysis of linear, and nonlinear, operators on cones both in finite and infinite dimensions, see [12, 13, 20]. In mathematical analysis one often works with Birkhoff's version of Hilbert's metric, which provides a slightly more general setup than the one outlined above. Birkhoff's version of Hilbert's metric, denoted d_H , is a metric on the set of rays in the interior, C° , of a closed cone C in a normed space X . If there exists a linear functional φ on X with $\varphi(x) > 0$ for all $x \in C \setminus \{0\}$, then δ_H and d_H coincide on $\Sigma_\varphi = \{x \in C^\circ : \varphi(x) > 0\}$, see for example [12, Theorem 2.1.2]. In general, however, there may not exist such a linear functional, see Remark 2.3 for more details, and in this respect Birkhoff's version is more general. Another advantage of using cones is that Hilbert's metric can be expressed in terms of the partial order induced by the cone, and one can use ideas from the theory of partially ordered vector spaces. In this paper we will be mainly working with Birkhoff's version of Hilbert's metric.

The paper has the following outline. In Sec. 2 we shall, besides introducing the relevant definitions, explain the relation between Hilbert's metric and Birkhoff's version of Hilbert's metric. Among other things we shall construct for a given Hilbert metric space (Ω, δ_H) in Y , a real normed vector space X containing Y such that on Ω the relative norm topology of X coincides with the Hilbert's metric topology.

Subsequently we prove in Sec. 3 the following theorem, which generalises [5, p. 163 (29.1)] and [7, Proposition 3].

Theorem 1.1. *If Ω_1, Ω_2 are strictly convex Hilbert geometries and $f : \Omega_1 \rightarrow \Omega_2$ is an isometry of (Ω_1, δ_H) into (Ω_2, δ_H) , then f is a projective linear homomorphism.*

The notion of a projective linear homomorphism will be given in Definition 2.7.

As we are working in infinite dimensions, one cannot use a projective basis and the fundamental theorem of projective geometry to prove Theorem 1.1. Instead we

establish an extension result, Proposition 3.3, which can be combined with Zorn’s Lemma to prove Theorem 1.1.

The second main result is proved in Sec. 4, and concerns the isometries of Hilbert geometries on infinite dimensional simplices. A natural generalisation of finite dimensional simplices to infinite dimensions is the set

$$\Delta(K, \mu)^\circ := \left\{ f \in C(K) : f(x) > 0 \text{ for all } x \in K \text{ and } \int f \, d\mu = 1 \right\},$$

where $C(K)$ is the set of continuous functions on a compact Hausdorff space K and μ is a finite, strictly positive, Borel measure on K , i.e. $\mu(K) < \infty$ and $\int f \, d\mu > 0$ for all $f \in C(K)$ with $f \neq 0$ and $f(x) \geq 0$ for all $x \in K$. Note that the set

$$\Delta_\infty^\circ := \left\{ x \in \ell^\infty : x_i > 0 \text{ for all } i \in \mathbb{N} \text{ and } \varphi(x) := \sum_{i=1}^\infty 2^{-i} x_i = 1 \right\}$$

is a special case. Indeed, as ℓ^∞ is an abstract M -space with an order unit, it follows from Kakutani’s representation theorem that ℓ^∞ is isometrically order-isomorphic to $C(K)$ for some compact Hausdorff space K , see [16, Theorem 1.b.6].

For the Hilbert geometries $(\Delta(K, \mu)^\circ, \delta_H)$ we have the following result.

Theorem 1.2. *If K_1 and K_2 are compact Hausdorff spaces with finite, strictly positive, Borel measures μ_1 and μ_2 , respectively, then $h : \Delta(K_1, \mu_1)^\circ \rightarrow \Delta(K_2, \mu_2)^\circ$ is a surjective Hilbert’s metric isometry if and only if there exist $\varepsilon \in \{-1, 1\}$, a homeomorphism $\vartheta : K_2 \rightarrow K_1$, and $g \in C(K_2)$ with $g(x) > 0$ for all $x \in K_2$ such that*

$$h(f) = \frac{g \cdot (f \circ \vartheta)^\varepsilon}{\int g \cdot (f \circ \vartheta)^\varepsilon \, d\mu_2}.$$

If $(K_1, \mu_1) = (K_2, \mu_2) = (K, \mu)$ and $|K| \geq 3$, then the isometry group is given by

$$\text{Isom}(\Delta(K, \mu)^\circ, \delta_H) \cong \overline{C(K)} \rtimes (C_2 \times \text{Homeo}(K)),$$

where C_2 is the cyclic group of order 2, $\overline{C(K)} := C(K)/\mathbb{R}\mathbf{1}$, and $\mathbf{1}$ is the constant one function on K .

Theorem 1.2 generalises the characterisation of the isometries of the Hilbert geometry on finite dimensional simplices obtained in [7], see also [8, Theorem 5.1] and [15, Theorem 1.2]. As a direct consequence we obtain the following result.

Corollary 1.3. *If K_1, K_2 are compact Hausdorff spaces, then $(\Delta(K_1, \mu_1)^\circ, d_H)$ and $(\Delta(K_2, \mu_2)^\circ, d_H)$ are isometric if and only if K_1 and K_2 are homeomorphic.*

2. Preliminaries

In this section we introduce the basic concepts and preliminary results. A cone C in a vector space X is a convex set such that $\lambda C \subseteq C$ for all $\lambda \geq 0$ and $C \cap -C = \{0\}$. A cone C induces a partial ordering \leq_C on X by $x \leq_C y$ if $y - x \in C$. The cone

is said to be *Archimedean* if for each $x \in X$ and $y \in C$ with $nx \leq_C y$ for all $n = 1, 2, 3, \dots$ we have that $x \leq_C 0$. An element $u \in C$ is called an *order unit* if for each $x \in X$ there exists $\lambda > 0$ such that $x \leq_C \lambda u$. The triple (X, C, u) is called an *order unit space*, if C is an Archimedean cone in X and u is an order unit for C .

Given an order unit space (X, C, u) , the space X can be equipped with the so-called *order unit norm*,

$$\|x\|_u := \inf\{\lambda > 0 : -\lambda u \leq_C x \leq_C \lambda u\}.$$

By [1, Theorem 2.55(2)], C is closed under $\|\cdot\|_u$. Note also that $\|\cdot\|_u$ is a *monotone norm*, i.e. $\|x\|_u \leq \|y\|_u$ for all $0 \leq_C x \leq_C y$, and hence C is normal with respect to $\|\cdot\|_u$. Recall that a cone C in a normed space $(X, \|\cdot\|)$ is called *normal* if there exists a constant $\kappa > 0$ such that $\|x\| \leq \kappa\|y\|$ whenever $0 \leq_C x \leq_C y$. Furthermore, C has nonempty interior, C° , with respect to $\|\cdot\|_u$, as the following lemma shows.

Lemma 2.1. *If (X, C, u) is an order unit space, then the set of order units of C coincides with C° .*

Proof. Every interior point is an order unit by [1, Lemma 2.5]. Conversely, if $x \in C$ is an order unit, then there exists $M > 0$ such that $u/M \leq x$. If $\|y\|_u \leq 1/M$, then $x - y \geq x - u/M \geq 0$, so $x \in C^\circ$. □

Throughout the paper we shall always assume that an order unit space is equipped with the order unit norm.

A linear functional $\varphi : X \rightarrow \mathbb{R}$ on an order unit space (X, C, u) is said to be *positive* if $\varphi(C) \subseteq [0, \infty)$. It is said to be *strictly positive* if $\varphi(C \setminus \{0\}) \subseteq (0, \infty)$. A positive linear functional φ with $\varphi(u) = 1$ is called a *state* of (X, C, u) . Strictly positive states are always continuous and $\|\varphi\| = 1$, as can be seen from the following lemma applied to $X_2 = \mathbb{R}$.

Lemma 2.2. *Let (X_1, C_1, u_1) and (X_2, C_2, u_2) be order unit spaces. If $T : X_1 \rightarrow X_2$ is a linear map such that $T(C_1) \subseteq C_2$, then T is continuous with $\|T\| = \|Tu_1\|_{u_2}$.*

Proof. If $\|x\|_{u_1} \leq 1$, then $-u_1 \leq_{C_1} x \leq_{C_1} u_1$, so that $-Tu_1 \leq_{C_2} Tx \leq_{C_2} Tu_1$. The statement now follows from the definition of $\|\cdot\|_{u_2}$. □

Remark 2.3. As mentioned in the Introduction, there may not exist a strictly positive functional for a given order unit space (X, C, u) . Indeed, if X is the vector space of bounded functions on an uncountable set K , C is the Archimedean cone of functions taking nonnegative values everywhere, and u is the constant one function on K , then no strictly positive functional exists, see [1, Exercise 6, Sec. 1.7]. However, if X is separable, then a strictly positive state always exists. Indeed, in that case the unit ball B_{X^*} of X^* (which is weak*-compact) is weak*-metrizable, and so there exists a sequence of states φ_n which is weak*-dense in the set of all states. A standard argument then shows that the state $\varphi := \sum_{n=1}^\infty 2^{-n}\varphi_n$ is strictly positive.

If (X, C, u) is an order unit space and $x, y \in C^\circ$, then it follows from Lemma 2.1 that there exist $0 < \alpha \leq \beta$ such that $\alpha y \leq_C x \leq_C \beta y$, and hence we can define

$$M(x/y) := \inf\{\beta > 0 : x \leq_C \beta y\} < \infty.$$

On C° (Birkhoff's version of) Hilbert's metric is given by

$$d_H(x, y) := \log(M(x/y)M(y/x)).$$

It is easy to verify that $d_H(\lambda x, \mu y) = d_H(x, y)$ for all $\lambda, \mu > 0$, and hence d_H is not a metric. However, it can be shown that d_H is a metric between pairs of rays in C° , see [13, Lemma 2.1]. We shall denote the projective spaces obtained by identifying points on rays inside C° by $P(C^\circ)$. So, $(P(C^\circ), d_H)$ is a metric space.

In case there exists a strictly positive state $\varphi : X \rightarrow \mathbb{R}$, we can identify $P(C^\circ)$ with the cross section

$$\Sigma_\varphi := \{x \in C^\circ : \varphi(x) = 1\}.$$

We shall use the following notation. For $x \in C^\circ$, we write $[x] := x/\varphi(x) \in \Sigma_\varphi$. The boundary of Σ_φ relative to the affine space $\{x \in X : \varphi(x) = 1\}$ in $(X, \|\cdot\|_u)$ is denoted by $\partial\Sigma_\varphi$. Its closure will be denoted by $\overline{\Sigma}_\varphi$.

The following theorem shows the relation between Hilbert geometries and order unit spaces with strictly positive functionals.

Theorem 2.4. *If Y is a vector space and $\Omega \subset Y$ is a convex set on which δ_H is well defined, then there exists an order unit space (X, C, u) and a strictly positive state φ on X such that Ω is affinely isomorphic to Σ_φ .*

Conversely, if (X, C, u) is an order unit space with a strictly positive state φ , then Σ_φ is a convex set on which δ_H is well defined.

Proof. Let $\Omega \subset Y$ be a convex set on which δ_H is well defined, i.e. $\Omega \cap \ell_{xy}$ is open and bounded for every $x, y \in \Omega$ with $x \neq y$. By translating we may assume without loss of generality that $0 \in \Omega$ and, by restricting to the span of Ω , we may also assume that $Y = \text{Span } \Omega$.

We claim that these assumptions imply that $\Omega \cap \ell_{0y}$ is open and bounded for every nonzero $y \in Y$. Indeed, let $y \in Y$ be nonzero, then $y = \sum_{i=1}^n \lambda_i y_i$ for some $y_i \in \Omega$ and $\lambda_i \in \mathbb{R}$, since Ω spans Y . In this representation we may assume that each $\lambda_i > 0$. Indeed, if $\lambda_i < 0$ for some i , then we can replace y_i by $-\varepsilon y_i$ for some small ε (since $\ell_{0y_i} \cap \Omega$ is open) and λ_i by $-\varepsilon^{-1} \lambda_i > 0$. Now if $\lambda := \sum_{i=1}^n \lambda_i$, then each $\lambda^{-1} \lambda_i > 0$ and they sum to 1. So,

$$\lambda^{-1}y = \sum_{i=1}^n \lambda^{-1} \lambda_i y_i \in \Omega$$

by convexity of Ω , and hence $\Omega \cap \ell_{0y} = \Omega \cap \ell_{0(\lambda^{-1}y)}$ is open and bounded.

By considering lines through 0, this implies that Ω is absorbing, i.e. for every $y \in Y$, there exists an $\varepsilon > 0$ such that $\lambda y \in \Omega$ for all $|\lambda| < \varepsilon$. Define

$$\overline{\Omega} := \{y \in Y : \lambda y \in \Omega \text{ for all } 0 \leq \lambda < 1\}$$

and note that $\overline{\Omega}$ is convex.

Let $X := Y \oplus \mathbb{R}$ and consider the cone

$$C_\Omega := \{\lambda(y, 1) \in X : \lambda \geq 0 \text{ and } y \in \overline{\Omega}\}.$$

We now show that C_Ω is Archimedean and $(0, 1) \in X$ is an order unit.

Let $(y, \lambda) \in X$ with $y \in Y$ and $\lambda \in \mathbb{R}$. To show that $(0, 1) \in X$ is an order unit, we have to find $M > 0$ with $-M(0, 1) \leq_{C_\Omega} (y, \lambda) \leq_{C_\Omega} M(0, 1)$. These inequalities hold if and only if $(y, \lambda + M), (-y, M - \lambda) \in C_\Omega$, which is equivalent to $y/(M + \lambda), -y/(\lambda - M) \in \overline{\Omega}$. But the existence of such an M now follows from the fact that $\overline{\Omega}$ is absorbing.

To show that C_Ω is Archimedean, it suffices to prove that $n(y, \lambda) \leq_{C_\Omega} (0, 1)$ for all large enough n implies that $-(y, \lambda) \in C_\Omega$, as $(0, 1)$ is an order unit. The assumed inequality is equivalent with $(-ny, 1 - n\lambda) \in C_\Omega$, which shows that $\lambda \leq 0$. If $\lambda = 0$, then $-ny \in \overline{\Omega}$, which implies by the boundedness of $\ell_{0y} \cap \Omega$ that $y = 0$ and so $-(y, \lambda) = (0, 0) \in C_\Omega$. If $\lambda < 0$, then by scaling we may assume that $\lambda = -1$, and then dividing the inequality by $1 + n$ yields

$$\frac{n}{1+n}(-y) \in \overline{\Omega}.$$

Hence $-y \in \overline{\Omega}$, and so $-(y, \lambda) = (-y, 1) \in C_\Omega$.

Obviously, the linear functional $\varphi : X \rightarrow \mathbb{R}$ defined by $\varphi((y, s)) := s$ is strictly positive with respect to C_Ω and $\varphi((0, 1)) = 1$, so φ is a strictly positive state.

Let us now show that Ω is affinely isomorphic to $\Sigma_\varphi := \{x \in C_\Omega^\circ : \varphi(x) = 1\}$. First assume that $y \in \overline{\Omega} \setminus \Omega$. Note that if we can prove that $(y, 1)$ is not an order unit, then $(y, 1) \notin C_\Omega^\circ$ by Lemma 2.1. Suppose by way of contradiction that $(y, 1)$ is an order unit. Then there exists $M > 1$ such that $(0, 1) \leq_{C_\Omega} M(y, 1)$, which is equivalent with $\frac{M}{M-1}y \in \overline{\Omega}$. Taking $0 < \lambda := \frac{M-1}{M} < 1$ we get that $y = \lambda \frac{M}{M-1}y \in \Omega$, which is absurd. Now suppose that $y \in \Omega$. We need to show that $(y, 1) \in C_\Omega^\circ$. As $(0, 1) \in C_\Omega^\circ$ is an order unit, it suffices to show by Lemma 2.1 that there exists $M > 1$ such that $(0, 1) \leq_{C_\Omega} M(y, 1)$. Recall that $\ell_{0y} \cap \Omega$ is an open subset of ℓ_{0y} . Hence there exists $M > 1$ such that $\frac{M}{M-1}y \in \ell_{0y} \cap \Omega$, which implies that $(0, 1) \leq_{C_\Omega} M(y, 1)$. We conclude that Ω is affinely isomorphic to $\Sigma_\varphi := \{x \in C_\Omega^\circ : \varphi(x) = 1\}$.

To prove the second part, we note that for distinct $x, y \in \Sigma_\varphi$ the points $w_x := x - M(y/x)^{-1}y$ and $w_y := y - M(x/y)^{-1}x$ are in $\partial C \setminus \{0\}$, as C is closed. So, $[w_x]$ and $[w_y]$ are the end-points of the straight line segment $\ell_{xy} \cap \Sigma_\varphi$. □

Remark 2.5. In the proof of Theorem 2.4, the vector space Y is a subspace of X , and so it inherits the norm $\|\cdot\|_u$ from X . Thus, for $y \in Y$,

$$\|y\| = \inf\{\lambda > 0 : -\lambda(0, 1) \leq (y, 0) \leq \lambda(0, 1)\}.$$

The condition on λ is equivalent with $(y, \lambda), (-y, \lambda) \in C_\Omega$, which in turn is equivalent with $y/\lambda, -y/\lambda \in \overline{\Omega}$, and so

$$\|y\| = \inf\{\lambda > 0 : y \in \lambda(\overline{\Omega} \cap -\overline{\Omega})\}.$$

Hence this norm equals the Minkowski functional of $\overline{\Omega} \cap -\overline{\Omega}$.

The advantage of working with cones is that we can use d_H instead of δ_H and apply ideas from the theory of partially ordered vector spaces. Indeed, the following result, which goes back to Birkhoff [2], is well known, see for example [12, Theorem 2.12 and Corollary 2.5.6].

Lemma 2.6. *If (X, C, u) is an order unit space with strictly positive state φ , then on Σ_φ the metrics d_H and δ_H coincide. Moreover, the Hilbert's metric topology on Σ_φ coincides with the order unit norm topology on X .*

If (X_1, C_1, u_1) and (X_2, C_2, u_2) are two order unit spaces and $T : X_1 \rightarrow X_2$ is a linear map, we say that T is *bi-positive* if $Tx \in C_2$ if and only if $x \in C_1$. Note that a bi-positive linear map $T : X_1 \rightarrow X_2$ is always injective, as $Tx = 0$ implies that x and $-x$ in C_1 , so that $x = 0$. Also note that if there exists $x \in C_1^\circ$ such that $Tx \in C_2^\circ$, then $T(C_1^\circ) \subseteq C_2^\circ$. Indeed, if $y \in C_1^\circ$, then y is an order unit by Lemma 2.1, and hence there exists $\lambda > 0$ such that $x \leq_{C_1} \lambda y$. It follows that $Tx \leq_{C_2} \lambda Ty$, so that Ty is an order unit, as Tx is an order unit, and hence $Ty \in C_2^\circ$.

It is easy to check that a bi-positive linear map $T : X_1 \rightarrow X_2$ induces a Hilbert's metric isometry, as $M(Tx/Ty) = M(x/y)$ for all $x, y \in C_1^\circ$. In that case we shall denote the induced map between the projective spaces $P(C_1^\circ)$ and $P(C_2^\circ)$ by $[T]$.

Definition 2.7. An isometry f from $(P(C_1^\circ), d_H)$ into $(P(C_2^\circ), d_H)$ is called a *projective linear homomorphism* if there exists a bi-positive linear map $T : X_1 \rightarrow X_2$ such that $f = [T]$ on $P(C_1^\circ)$.

Before we start the proof of Theorem 1.1, we collect some final pieces of notation. Recall that the image of a map γ from a (possibly unbounded) interval $I \subseteq \mathbb{R}$ into (Σ_φ, d_H) is *geodesic* if

$$d_H(\gamma(t), \gamma(s)) = |t - s| \quad \text{for all } t, s \in I.$$

Given a straight line ℓ_{xy} through $x \neq y$ in Σ_φ , we write $\ell_{xy}^+ := \ell_{xy} \cap \Sigma_\varphi$. Also for $x, y \in X$ the closed and open line segments are, respectively, denoted by

$$[x, y] := \{tx + (1 - t)y : 0 \leq t \leq 1\} \quad \text{and} \quad (x, y) := \{tx + (1 - t)y : 0 < t < 1\}.$$

The half-open intervals are defined in a similar way. For each $x, y \in \Sigma_\varphi$ the segment $[x, y]$ is geodesic in (Σ_φ, d_H) , see [10]. It is, however, in general not the only geodesic. The unique geodesics in (Σ_φ, d_H) are characterised as follows, see [10] or [7, Proposition 2].

Lemma 2.8. *Let (X, C, u) be an order unit space with strictly positive state φ . If $x, y \in \Sigma_\varphi$ and $x', y' \in \partial\Sigma_\varphi$ are the end points of $\ell_{xy} \cap \Sigma_\varphi$, then $[x, y]$ is the*

unique geodesic connecting x and y in (Σ_φ, d_H) if and only if there exist no open line segments $I_{x'}$ through x' and $I_{y'}$ through y' in $\partial\Sigma_\varphi$ such that the affine span of $I_{x'} \cup I_{y'}$ is 2-dimensional.

3. Strictly Convex Hilbert Geometries

We prove Theorem 1.1. Throughout this section we shall assume that (X_1, C_1, u_1) and (X_2, C_2, u_2) are order unit spaces, with strictly positive states $\varphi_i : X_i \rightarrow \mathbb{R}$ for $i = 1, 2$. For simplicity we write

$$\Sigma_i := \{x \in C_i^\circ : \varphi_i(x) = 1\} \quad \text{and} \quad \|\cdot\|_i := \|\cdot\|_{u_i} \quad \text{for } i = 1, 2.$$

Recall that Σ_i , $i = 1, 2$, is *strictly convex* if for each $x, y \in \partial\Sigma_i$ we have that $(x, y) \subseteq \Sigma_i$. In that case, it follows from Lemma 2.8 that the metric spaces (Σ_i, d_H) , $i = 1, 2$, are uniquely geodesic. Thus, any isometry f of (Σ_1, d_H) into (Σ_2, d_H) has to map line segments to line segments if (Σ_1, d_H) and (Σ_2, d_H) are strictly convex. We shall show that any isometry that maps line segments to line segments must be a projective linear homomorphism, which implies Theorem 1.1. We begin with the following lemma, which generalises [7, p. 101].

Lemma 3.1. *If $f : (\Sigma_1, d_H) \rightarrow (\Sigma_2, d_H)$ is an isometry that maps line segments to line segments, then f has a unique continuous extension to a map from $(\overline{\Sigma}_1, \|\cdot\|_1)$ into $(\overline{\Sigma}_2, \|\cdot\|_2)$. Furthermore, this extension is injective.*

Proof. The uniqueness of the continuous extension follows from the density of Σ_1 in $\overline{\Sigma}_1$. Convergence in this proof will always be in the order unit norm; recall that on Σ_i , the order unit norm topology and d_H -topology coincide by Lemma 2.6.

Fix an element $p \in \Sigma_1$ and let $x \in \partial\Sigma_1$. The line segment $[p, x]$ is mapped onto the line segment $[f(p), \xi]$, for some $\xi \in \partial\Sigma_2$, because

$$d_H(p, (1 - t)p + tx) \rightarrow \infty$$

as $t \uparrow 1$. We define $f(x) := \xi$.

To show continuity of the extension, let $(x_n)_n$ be a sequence in $\overline{\Sigma}_1$ converging to x . Then $y_n := (p + x_n)/2 \rightarrow (p + x)/2 =: y$, and so $f(y_n) \rightarrow f(y)$. Let $s > 1$ be such that $sf(y) + (1 - s)f(p) \notin \overline{\Sigma}_2$. Then for some $N \geq 1$ and all $n \geq N$,

$$sf(y_n) + (1 - s)f(p) \notin \overline{\Sigma}_2. \tag{3.1}$$

Since f maps line segments to line segments, there exist s_n such that

$$s_n f(y_n) + (1 - s_n)f(p) = f(x_n) \in \overline{\Sigma}_2,$$

and combining this with (3.1) yields $s_n < s$ for all $n \geq N$.

Now suppose that $f(x_n)$ does not converge to $f(x)$. By passing to a subsequence three times, we find a subsequence $(x_{n_k})_k$ such that $f(x_{n_k})$ stays away from $f(x)$, $s_{n_k} \rightarrow r \in [0, s]$, and that either all $x_{n_k} \in \partial\Sigma_1$ or all $x_{n_k} \in \Sigma_1$. It follows that

$$f(x_{n_k}) = s_{n_k} f(y_{n_k}) + (1 - s_{n_k})f(p) \rightarrow rf(y) + (1 - r)f(p) \in \overline{\Sigma}_2.$$

We claim that $rf(y) + (1-r)f(p) \in \partial\Sigma_2$. Indeed, if all $x_{n_k} \in \partial\Sigma_1$, then $f(x_{n_k}) \in \partial\Sigma_2$ and the claim follows from the closedness of $\partial\Sigma_2$. If all $x_{n_k} \in \Sigma_1$, then the fact f is a d_H -isometry combined with $d_H(x_{n_k}, p) \rightarrow \infty$ yields $d_H(f(x_{n_k}), f(p)) \rightarrow \infty$, and so $rf(y) + (1-r)f(p) \in \partial\Sigma_2$.

Hence $f(x) = rf(y) + (1-r)f(p)$ by construction of $f(x)$, and so $f(x_{n_k}) \rightarrow f(x)$, which is impossible since $f(x_{n_k})$ stays away from $f(x)$. Therefore the extension of f is continuous.

To show injectivity suppose that $x, y \in \bar{\Sigma}_1$ are such that $x \neq y$. Note that $[f(p), f(x)] \neq [f(p), f(y)]$, since f is injective on Σ_1 ; so, $f(x) \neq f(y)$. □

The following lemma is essentially part (iii) of the lemma on p. 101 in [7], and will be useful in the sequel.

Lemma 3.2. *Let $x, y \in \Sigma_1$ and let $x', y' \in \partial\Sigma_1$ be the end points of ℓ_{xy}^+ such that x is between x' and y , and y is between y' and x . Suppose that $f : (\Sigma_1, d_H) \rightarrow (\Sigma_2, d_H)$ is an isometry that maps line segments to line segments. If $f(x)'$ and $f(y)'$ are the end points of $\ell_{f(x)f(y)}^+$ such that $f(x)$ is between $f(x)'$ and $f(y)$, and $f(y)$ is between $f(y)'$ and $f(x)$, then there exists a linear map $S : \text{Span}\{x', y'\} \rightarrow \text{Span}\{f(x)', f(y)'\}$ satisfying $[Sx'] = f(x)'$, $[Sy'] = f(y)'$, $[Sz] = f(z)$ for all $z \in (x', y')$, and S is bi-positive with respect to $C_1 \cap \text{Span}\{x', y'\}$ and $C_2 \cap \text{Span}\{f(x)', f(y)'\}$.*

Proof. Let $0 < s, t < 1$ be such that $x = tx' + (1-t)y'$ and $f(x) = sf(x') + (1-s)f(y')$. Since $\{tx', (1-t)y'\}$ and $\{sf(x'), (1-s)f(y')\}$ define bases for $\text{Span}\{x', y'\}$ and $\text{Span}\{f(x)', f(y)'\}$, and the cones $C_1 \cap \text{Span}\{x', y'\}$ and $C_2 \cap \text{Span}\{f(x)', f(y)'\}$ are the positive span of these basis elements, we have a bijective and bi-positive linear map $S : \text{Span}\{x', y'\} \rightarrow \text{Span}\{f(x)', f(y)'\}$ defined by

$$S(\alpha x' + \beta(1-t)y') := \alpha sf(x') + \beta(1-s)f(y)'$$

Note that $[Sx'] = f(x)'$, $[Sy'] = f(y)'$ and $Sx = f(x)$.

Let $z \in [x, x')$ and note that, as f maps line segments to line segments and f is an isometry, we have that

$$[f(x)', f(z), f(x), f(y)'] = \exp(d_H(f(z), f(x))) = \exp(d_H(z, x)) = [x', z, x, y'].$$

As the cross ratio is a projective invariant, we know that

$$[[Sx'], [Sz], [Sx], [Sy']] = [Sx', Sx, Sy, Sy'] = [x', z, x, y'],$$

which combined with the previous equality gives

$$[[Sx'], [Sz], [Sx], [Sy']] = [f(x)', f(z), f(x), f(y)'] = [[Sx'], f(z), [Sx], [Sy']].$$

This implies that $[Sz] = f(z)$. Interchanging the roles of x' and y' finally gives $[Sz] = f(z)$ for all $z \in (x', y')$. □

Proposition 3.3. *Let $f : (\Sigma_1, d_H) \rightarrow (\Sigma_2, d_H)$ be an isometry that maps line segments to line segments and $Y \subseteq X_1$ be a subspace such that $Y \cap \Sigma_1 \neq \emptyset$. If*

$T : Y \rightarrow X_2$ is a bi-positive linear map such that $[Ty] = f(y)$ for all $y \in Y \cap \Sigma_1$, then for $z \in \Sigma_1 \setminus Y$ there is a bi-positive linear extension

$$\hat{T} : Y \oplus \text{Span}\{z\} \rightarrow X_2$$

of T such that $[\hat{T}y] = f(y)$ for all $y \in (Y \oplus \text{Span}\{z\}) \cap \Sigma_1$.

Proof. Let $T : Y \rightarrow X_2$ be a bi-positive linear map that satisfies $[Ty] = f(y)$ for all $y \in Y \cap \Sigma_1$ and choose $z \in \Sigma_1 \setminus Y$. Fix $\xi \in Y \cap \Sigma_1$ and consider $\ell_{z\xi}^+$. By Lemma 3.2 there exists a bi-positive linear map $S : \text{Span}\{\xi, z\} \rightarrow X_2$ such that $[Sx] = f(x)$ for all $x \in \ell_{z\xi}^+$. By rescaling S , we may assume that $S\xi = T\xi$.

Now let $\hat{Y} := Y \oplus \text{Span}\{z\}$ and define the linear map

$$\hat{T} : \hat{Y} \rightarrow \text{ran } T + \text{Span}\{Sz\}$$

by $y + \lambda z \mapsto Ty + \lambda Sz$. We wish to show that $\text{ran } T + \text{Span}\{Sz\}$ is in fact a direct sum, making \hat{T} injective. Suppose $Ty = \lambda Sz$ for some $y \in Y$ and $\lambda > 0$. As T is bi-positive, this implies that $y \in Y \cap C_1^\circ$, so $[Ty] = [Sz] = f(z)$. But this yields $f([y]) = f(z)$, so $[y] = z$ and this is impossible. If $\lambda < 0$, we can use a similar argument for $-y$ to arrive at a contradiction. Thus, $\text{ran } T + \text{Span}\{Sz\}$ is indeed a direct sum. Since $\hat{T}\xi = T\xi = S\xi$ and $\hat{T}z = Sz$, it follows that $\hat{T} = S$ on $\ell_{z\xi}^+$ and so $[\hat{T}x] = f(x)$ for all $x \in \ell_{z\xi}^+$.

Now suppose that $w \in \hat{Y} \cap \Sigma_1$ and $w \notin Y \cup \ell_{z\xi}^+$. The subspace Y is a hyperplane in \hat{Y} and therefore, it divides $\hat{Y} \cap \Sigma_1$ into two parts. Choose distinct $\eta_1, \eta_2 \in \ell_{z\xi}^+$ that lie on the other side of w and let $y_1, y_2 \in Y \cap \Sigma_1$ be the intersection points of the line segments $\ell_{w\eta_1}^+$ and $\ell_{w\eta_2}^+$, respectively. The situation is depicted in Fig. 1 below. We see that w is the unique point of intersection of the line segments $\ell_{\eta_1 y_1}^+$ and $\ell_{\eta_2 y_2}^+$. As f is injective, it follows that $f(w)$ is the unique point of intersection of the line segments $\ell_{f(\eta_1) f(y_1)}^+$ and $\ell_{f(\eta_2) f(y_2)}^+$. Also, as $[\hat{T}\eta_i] = f(\eta_i)$ and $[\hat{T}y_i] = f(y_i)$, we have $\hat{T}\eta_i, \hat{T}y_i \in \text{Span}\{f(\eta_i), f(y_i)\}$; hence

$$\hat{T}w \in \text{Span}\{f(\eta_1), f(y_1)\} \cap \text{Span}\{f(\eta_2), f(y_2)\} = \text{Span}\{f(w)\}.$$




Fig. 1. Points of intersection.

So, $\hat{T}w \in C_2^\circ \cup -C_2^\circ$, since \hat{T} is injective, and hence

$$\hat{T}(\hat{Y} \cap C_1^\circ) \subseteq C_2^\circ \cup -C_2^\circ.$$

The convexity of $\hat{Y} \cap C_1^\circ$ now implies that either $\hat{T}(\hat{Y} \cap C_1^\circ) \subseteq C_2^\circ$ or $\hat{T}(\hat{Y} \cap C_1^\circ) \subseteq -C_2^\circ$. As $\hat{T}\xi = T\xi \in C_2^\circ$, it follows that $\hat{T}(\hat{Y} \cap C_1^\circ) \subseteq C_2^\circ$. Moreover, this shows that $[\hat{T}w]$ is well defined. As w was arbitrary, we conclude that $[\hat{T}y] = f(y)$ for all $y \in \hat{Y} \cap \Sigma_1$.

Next, we will show that \hat{T} is bi-positive. Suppose $x \in \partial C_1 \cap \hat{Y}$ with $x \neq 0$. We claim that $\hat{T}(x) \in \partial C_2$. Since $\varphi_1(x) > 0$, we may assume without loss of generality that $x \in \partial \Sigma_1$. Define $x_n := (1 - \frac{1}{n})x + \frac{1}{n}\xi$ for $n \geq 1$. Then we have $x_n \rightarrow x$, so that $f(x_n) \rightarrow f(x)$ by Lemma 3.1 and

$$\hat{T}x_n = \hat{T}|_{\text{Span}\{x,\xi\}}x_n \rightarrow \hat{T}|_{\text{Span}\{x,\xi\}}x = \hat{T}x$$

as $\dim(\text{Span}\{x, \xi\}) < \infty$. It follows that $\varphi_2(\hat{T}x_n) \rightarrow \varphi_2(\hat{T}x)$ and, since $\hat{T}x_n \in C_2$ for all $n \geq 1$, we must have $\hat{T}x \in C_2$, because C_2 is closed. Moreover, the injectivity of \hat{T} yields $\varphi_2(\hat{T}x) > 0$. We also have that

$$\hat{T}x_n = \varphi_2(\hat{T}x_n)f(x_n) \rightarrow \varphi_2(\hat{T}x)f(x) \in \partial C_2,$$

so we conclude that $\hat{T}x \in \partial C_2$ showing that \hat{T} is positive. Finally, if we pick $x \in \hat{Y} \setminus C_1$, then there exists a $0 < t < 1$ such that $tx + (1 - t)\xi \in \partial C_1$ and so $t\hat{T}x + (1 - t)\hat{T}\xi \in \partial C_2$ by our previous findings. But this implies that $\hat{T}x \notin C_2$ making \hat{T} bi-positive. □

Enough preparations have been made to prove the following result which implies Theorem 1.1 by Theorem 2.4.

Theorem 3.4. *An map $f : (\Sigma_1, d_H) \rightarrow (\Sigma_2, d_H)$ is an isometry that maps line segments to line segments if and only if it is a projective linear homomorphism.*

Proof. Consider the collection \mathcal{C} of pairs (Y, T_Y) where $Y \subseteq X_1$ is a linear subspace and $T_Y : Y \rightarrow X_2$ is a bi-positive linear map such that $[T_Y x] = f(x)$ for all $x \in Y \cap \Sigma_1$. Note that $\mathcal{C} \neq \emptyset$, since $C_1^\circ \neq \emptyset$. We can define a partial order \leq on \mathcal{C} by

$$(Y, T_Y) \leq (Z, T_Z) \quad \text{if } Y \subseteq Z \text{ and } T_Z y = T_Y y \text{ for all } y \in Y.$$

Let $(Y_i, T_{Y_i})_{i \in I}$ be a totally ordered subset in \mathcal{C} . Put $Y := \bigcup_{i \in I} Y_i$ and define $T_Y : Y \rightarrow X_2$ through $T_Y y_i := T_{Y_i} y_i$. Clearly, we have that Y is a linear subspace of X_1 and T_Y is a well-defined bi-positive linear map. For $y \in Y \cap \Sigma_1$ we have $y \in Y_i \cap \Sigma_1$ for some $i \in I$ and $[T_Y y] = [T_{Y_i} y] = f(y)$, so $(Y, T_Y) \in \mathcal{C}$ is an upper bound. By Zorn's lemma our collection \mathcal{C} contains a maximal element (Ω, T_Ω) . Suppose that $x \in \Sigma_1 \setminus \Omega$. By Proposition 3.3 we have a bi-positive linear extension

$$\hat{T}_\Omega : \Omega \oplus \text{Span}\{x\} \rightarrow X_2$$

of T_Ω such that $[\hat{T}_\Omega z] = f(z)$ for all $z \in (\Omega \oplus \text{Span}\{x\}) \cap \Sigma_1$. But now we have $(\Omega \oplus \text{Span}\{x\}, \hat{T}_\Omega) \in \mathcal{C}$ and $(\Omega, T_\Omega) \leq (\Omega \oplus \text{Span}\{x\}, \hat{T}_\Omega)$, which contradicts the

maximality of (Ω, T_Ω) . We conclude that $\Omega \cap \Sigma_1 = \Sigma_1$ and therefore $[T_\Omega x] = f(x)$ for all $x \in \Sigma_1$.

We claim that $\Omega = X_1$. Let $x \in X_1, u \in \Sigma_1$ and $\varepsilon > 0$ such that $B_\varepsilon(u) \subseteq C_1^\circ$. Also, there are $y, z \in C_1$ such that $x = y - z$, as $C_1 - C_1 = X_1$. Now, for $k \geq 1$ large enough, we have that $\frac{1}{k}y + u, \frac{1}{k}z + u \in B_\varepsilon(u)$. Since $\Sigma_1 \subseteq \Omega$, it follows that $C_1^\circ \subseteq \Omega$; hence

$$\frac{1}{k}x = \left(\frac{1}{k}y + u\right) - \left(\frac{1}{k}z + u\right) \in \Omega,$$

so $x \in \Omega$.

Obviously any projective linear homomorphism maps line segments to line segments, and hence we are done. □

Note that Lemma 2.2 shows that the linear map T in the previous theorem is continuous with respect to the order unit norm. We also note that Theorem 3.4 implies that two uniquely geodesic Hilbert geometries Ω_1 and Ω_2 are isometric if and only if there exists a projective linear isomorphism between them.

Remark 3.5. An important variant of Hilbert’s metric is Thompson’s metric which was introduced in [23]. On the interior of a cone C in an order unit space, *Thompson’s metric* is given by

$$d_T(x, y) := \log \max\{M(x/y), M(y/x)\} \quad \text{for } x, y \in C^\circ.$$

It was shown in [14, Theorem 8.2] that if C is a finite dimensional strictly convex cone with $\dim C \geq 3$, then for every isometry f of (C°, d_T) there exists a bi-positive linear map $T : X \rightarrow X$ such that for each $x \in C^\circ$ we have that $f(x) = \lambda_x T x$ for some $\lambda_x > 0$. The proof of this result relies on [7, Proposition 3]. Using Theorem 3.4 it is straightforward to extend [14, Theorem 8.2] to infinite dimensional strictly convex cones.

4. Infinite Dimensional Simplices

Let K be a compact Hausdorff space and $C(K)$ denote the space of real-valued continuous functions on K . Consider the cone $C(K)_+$ consisting of nonnegative functions with interior,

$$C(K)_+^\circ := \{f \in C(K) : f(x) > 0 \text{ for all } x \in K\}.$$

It is well known that the Hilbert geometry on a finite dimensional simplex is isometric to a finite dimensional normed space. The same is true for $(P(C(K)_+^\circ), d_H)$, see [20, Proposition 1.7]. It will be useful to recall the basic argument. Let $\mathbf{1}$ be the constant one function on K and denote the elements in the quotient space $\overline{C(K)} := C(K)/\mathbb{R}\mathbf{1}$ by \bar{g} . The map $\text{Log} : P(C(K)_+^\circ) \rightarrow \overline{C(K)}$ given by,

$$\text{Log}(\bar{f}) := \overline{\log \circ f} \quad \text{for } \bar{f} \in P(C(K)_+^\circ),$$

is an isometry of $(P(C(K))_+^\circ, d_H)$ onto $(\overline{C(K)}, \|\cdot\|_{\text{var}})$, where

$$\|\overline{g}\|_{\text{var}} := \sup_{x \in K} g(x) - \inf_{x \in K} g(x),$$

is the *variation norm*. To see this note that

$$M(f/g) = \inf\{\beta > 0 : f(x) \leq \beta g(x) \text{ for all } x \in K\} = \sup_{x \in K} \frac{f(x)}{g(x)}$$

for $f, g \in P(C(K)_+^\circ)$, so that

$$\begin{aligned} d_H(f, g) &= \sup_{x \in K} \frac{\log f(x)}{\log g(x)} + \sup_{x \in K} \frac{\log g(x)}{\log f(x)} \\ &= \sup_{x \in K} (\log f(x) - \log g(x)) - \inf_{x \in K} (\log f(x) - \log g(x)) \\ &= \|\text{Log}(\overline{f}) - \text{Log}(\overline{g})\|_{\text{var}}. \end{aligned}$$

Given a function $f \in C(K)$ the supremum norm of its translations $f - \lambda \mathbf{1}$ for $\lambda \in \mathbb{R}$ is minimised precisely when translating f by the average of both extreme values $\frac{1}{2}(\sup_{x \in K} f(x) + \inf_{x \in K} f(x))$. This means that the quotient norm on $C(K)/\mathbb{R}\mathbf{1}$ with respect to the supremum norm is exactly half of $\|\cdot\|_{\text{var}}$. This assertion is made precise in the following lemma.

Lemma 4.1. *Let K be a compact Hausdorff space. If $\|\cdot\|_q$ is the quotient norm on $\overline{C(K)}$ with respect to $2\|\cdot\|_\infty$, then $\|\cdot\|_q$ coincides with $\|\cdot\|_{\text{var}}$ on $\overline{C(K)}$.*

Proof. Let $f \in C(K)$ and write $a \vee b = \max\{a, b\}$ for $a, b \in \mathbb{R}$. Using the elementary fact that $\inf_{\lambda \in \mathbb{R}} (a - \lambda) \vee (b + \lambda) = (a + b)/2$, we see that

$$\begin{aligned} \|\overline{f}\|_q &= 2 \inf_{\lambda \in \mathbb{R}} \|f - \lambda \mathbf{1}\|_\infty = 2 \inf_{\lambda \in \mathbb{R}} \sup_{s \in K} |f(s) - \lambda| \\ &= 2 \inf_{\lambda \in \mathbb{R}} \left[\left(\sup_{s \in K} f(s) - \lambda \right) \vee \left(\sup_{s \in K} -f(s) + \lambda \right) \right] \\ &= \sup_{s \in K} f(s) + \sup_{s \in K} -f(s) = \sup_{s \in K} f(s) - \inf_{s \in K} f(s) = \|\overline{f}\|_{\text{var}}. \quad \square \end{aligned}$$

Our findings so far have shown that in order to describe the surjective isometries

$$h : (P(C(K_1)_+^\circ), d_H) \rightarrow (P(C(K_2)_+^\circ), d_H),$$

it suffices to understand the surjective isometries

$$T : (\overline{C(K_1)}, \|\cdot\|_q) \rightarrow (\overline{C(K_2)}, \|\cdot\|_q).$$

By the Mazur–Ulam theorem these isometries T must be affine. We can compose T with an appropriate translation to make it linear. Thus, our goal will be to classify all isometric isomorphisms (surjective linear isometries) $T : (\overline{C(K_1)}, \|\cdot\|_q) \rightarrow (\overline{C(K_2)}, \|\cdot\|_q)$.

We will follow the lines of the proof for the Banach–Stone theorem [6, Theorem VI.2.1]. The Banach–Stone theorem characterises the isometric isomorphisms between $(C(K), \|\cdot\|_\infty)$ spaces. A common way to prove the Banach–Stone theorem is by looking at the adjoint operator, which is an isometry on the dual of $(C(K), \|\cdot\|_\infty)$, and to exploit the extreme points of the unit ball there. We shall take a similar approach.

The dual of $(C(K), \|\cdot\|_\infty)$ is $(M(K), \|\cdot\|_{TV})$, where $M(K)$ is the space of all regular signed Borel measures on K and $\|\mu\|_{TV} := |\mu|(K)$ is the *total variation norm*. Let us recall some basic facts about $(M(K), \|\cdot\|_{TV})$, which can be found in [6, Appendix C].

Every $\mu \in M(K)$ has a Hahn–Jordan decomposition $\mu = \mu^+ - \mu^-$ where μ^+ and μ^- are positive measures in $M(K)$, and

$$\|\mu\|_{TV} = \|\mu^+\|_{TV} + \|\mu^-\|_{TV}. \tag{4.1}$$

Also, if $\mu, \nu \in M(K)$ are positive, we have $\|\mu + \nu\|_{TV} = \|\mu\|_{TV} + \|\nu\|_{TV}$. The space $M(K)$ is a lattice: every $\mu, \nu \in M(K)$ have a supremum (least upper bound) $\mu \vee \nu$. The set $P(K)$ denotes the set of probability measures on K , and its extreme points are the set of Dirac measures $\{\delta_s : s \in K\}$. The map $s \mapsto \delta_s$ is a homeomorphism from K onto $\{\delta_s : s \in K\}$ equipped with the weak*-topology.

The dual space of $(C(K)/\mathbb{R}\mathbf{1}, \|\cdot\|_q)$ is $\mathbb{R}\mathbf{1}^\perp \subseteq (M(K), \frac{1}{2}\|\cdot\|_{TV})$, where $\mathbb{R}\mathbf{1}^\perp := \{\mu \in M(K) : \mu(K) = 0\}$. It follows that

$$\mathbb{R}\mathbf{1}^\perp = \{\mu \in M(K) : \|\mu^+\|_{TV} = \|\mu^-\|_{TV}\}.$$

Now, if

$$T : (\overline{C(K_1)}, \|\cdot\|_q) \rightarrow (\overline{C(K_2)}, \|\cdot\|_q)$$

is an isometric isomorphism, then the corresponding adjoint operator

$$T^* : \left(\mathbb{R}\mathbf{1}_2^\perp, \frac{1}{2}\|\cdot\|_{TV}\right) \rightarrow \left(\mathbb{R}\mathbf{1}_1^\perp, \frac{1}{2}\|\cdot\|_{TV}\right)$$

is an isometric isomorphism as well. Moreover, T^* is a weak*-homeomorphism from the unit ball $B_2 \subseteq \mathbb{R}\mathbf{1}_2^\perp$ onto the unit ball $B_1 \subseteq \mathbb{R}\mathbf{1}_1^\perp$ that maps the set of extreme points of B_2 , denoted $\text{ext}(B_2)$, bijectively onto the set of the extreme points of B_1 , denoted $\text{ext}(B_1)$. The following lemma tells us that the extreme points are exactly the differences of Dirac measures.

Proposition 4.2. *Let K be a compact Hausdorff space. The set of extreme points, $\text{ext}(B)$, of the unit sphere B in $\mathbb{R}\mathbf{1}^\perp \subseteq (M(K), \frac{1}{2}\|\cdot\|_{TV})$ satisfies*

$$\text{ext}(B) = \{\delta_s - \delta_t : s, t \in K \text{ and } s \neq t\}.$$

Proof. Let $\delta_s - \delta_t \in B$ with $s \neq t$ and suppose $\mu, \nu \in B$ are such that

$$\delta_s - \delta_t = \frac{1}{2}(\mu + \nu) = \frac{1}{2}(\mu^+ + \nu^+) - \frac{1}{2}(\mu^- + \nu^-).$$

Then $\delta_s = (\delta_s - \delta_t) \vee 0 \leq \frac{1}{2}(\mu^+ + \nu^+)$, and so for $\eta := \frac{1}{2}(\mu^+ + \nu^+) - \delta_s \geq 0$ it follows from (4.1) that

$$1 + \|\eta\|_{TV} = \|\delta_s + \eta\|_{TV} = \frac{1}{2}\|\mu^+ + \nu^+\|_{TV} = \frac{1}{2}\|\mu^+\|_{TV} + \frac{1}{2}\|\nu^+\|_{TV} = 1,$$

so $\eta = 0$, which yields $\delta_s = \frac{1}{2}(\mu^+ + \nu^+)$. The fact that δ_s is an extreme point in $B_{M(K)}$ (see [6, Theorem V.8.4]) implies that $\mu^+ = \nu^+$. Similarly, we have that $\delta_t \leq \frac{1}{2}(\mu^- + \nu^-)$ which implies $\mu^- = \nu^-$. This shows that $\delta_s - \delta_t \in \text{ext}(B)$.

Conversely, let $\mu = \mu^+ - \mu^- \in \text{ext}(B)$, and suppose $\mu^+ = \frac{1}{2}(\nu_1 + \nu_2)$ for $\nu_1, \nu_2 \in P(K)$. Then for $i = 1, 2$,

$$\frac{1}{2}\|\nu_i - \mu^-\|_{TV} \leq \frac{1}{2}(\|\nu_i\|_{TV} + \|\mu^-\|_{TV}) = 1,$$

so $\nu_i - \mu^- \in B$. Since $\mu \in \text{ext}(B)$ and $\mu = \frac{1}{2}(\nu_1 - \mu^-) + \frac{1}{2}(\nu_2 - \mu^-)$, we obtain $\nu_1 = \nu_2$. Hence $\mu^+ \in \text{ext}(P(K))$, which implies that $\mu^+ = \delta_s$ for some $s \in K$ by [6, Theorem V.8.4]. A similar argument yields $\mu^- = \delta_t$ for some $t \in K$. □

For $s \in K$ define the sets $E_s := \{\delta_s - \delta_t : t \in K, s \neq t\}$. Clearly the distance between distinct elements of E_s is 1, and it turns out that the sets $\pm E_s$ are the maximal equilateral subsets in $\text{ext}(B)$ of mutual distance 1.

Lemma 4.3. *Let K be a compact Hausdorff space and $\emptyset \neq A \subseteq \text{ext}(B)$ be such that $\frac{1}{2}\|\mu - \nu\|_{TV} = 1$ for all $\mu, \nu \in A$ with $\mu \neq \nu$, then there is an element $s \in K$ such that $A \subseteq E_s$ or $A \subseteq -E_s$.*

Proof. If such an s does not exist, then there exist elements $\delta_s - \delta_t, \delta_p - \delta_q \in A$ with $s \neq p$ and $t \neq q$. But clearly

$$\frac{1}{2}\|(\delta_s - \delta_t) - (\delta_p - \delta_q)\|_{TV} = 2. \quad \square$$

Now let $T : \overline{C(K_1)} \rightarrow \overline{C(K_2)}$ be an isometric isomorphism. Then the isometric isomorphism T^* preserves the maximal equilateral subsets of the extreme points of mutual distance 1. Hence $T^*(E_s) = \pm E_{\vartheta(s)}$. Note that if $s \neq t$, then $E_s \cap E_t = \emptyset$. But $E_s \cap -E_t \neq \emptyset$, as $\delta_s - \delta_t \in E_s \cap -E_t$. As T^* maps disjoint sets to disjoint sets, either $T^*(E_s) = E_{\vartheta(s)}$ for all $s \in K_2$, or, $T^*(E_s) = -E_{\vartheta(s)}$ for all $s \in K_2$. Thus, there exists $\varepsilon \in \{-1, 1\}$ such that $T^*(E_s) = \varepsilon E_{\vartheta(s)}$ for all $s \in K_2$, and ϑ is a bijection from K_2 to K_1 .

Lemma 4.4. *The above constructed bijection $\vartheta : K_2 \rightarrow K_1$ is a homeomorphism.*

Proof. Let $(s_\alpha)_\alpha$ be a net in K_2 converging to $s \in K_2$. Then δ_{s_α} converges weak* to δ_s , and so

$$\delta_{\vartheta(s_\alpha)} - \delta_{\vartheta(s)} = \varepsilon T^*(\delta_{s_\alpha} - \delta_s) \xrightarrow{\text{weak}^*} \varepsilon T^*0 = 0.$$

Hence $\delta_{\vartheta(s_\alpha)} \xrightarrow{\text{weak}^*} \delta_{\vartheta(s)}$, or equivalently, $\vartheta(s_\alpha) \rightarrow \vartheta(s)$. So ϑ is a continuous bijection from a compact space into a Hausdorff space and so it is a homeomorphism. □

We are now able to prove our main result concerning isometric isomorphisms $T : \overline{C(K_1)} \rightarrow \overline{C(K_2)}$.

Theorem 4.5. *If K_1, K_2 are compact Hausdorff spaces, then $T : \overline{C(K_1)} \rightarrow \overline{C(K_2)}$ is an isometric isomorphism if and only if there exist an $\varepsilon \in \{-1, 1\}$ and a homeomorphism $\vartheta : K_2 \rightarrow K_1$ such that $T\overline{f} = \varepsilon(\overline{f \circ \vartheta})$.*

Proof. Suppose T is an isometric isomorphism, and let ϑ and ε be such that $T^*E_s = \varepsilon E_{\vartheta(s)}$. Although point-evaluation for elements $\overline{g} \in \overline{C(K_2)}$ is not well-defined, the values $\overline{g}(s) - \overline{g}(t)$ for $s, t \in K_2$ are well-defined, and the computation

$$\begin{aligned} T\overline{f}(s) - T\overline{f}(t) &= (\delta_s - \delta_t)(T\overline{f}) \\ &= [T^*(\delta_s - \delta_t)](\overline{f}) \\ &= [\varepsilon(\delta_{\vartheta(s)} - \delta_{\vartheta(t)})](\overline{f}) \\ &= \varepsilon(\overline{f}(\vartheta(s)) - \overline{f}(\vartheta(t))) \end{aligned}$$

shows that T is induced from the map $f \mapsto \varepsilon(f \circ \vartheta)$ from $C(K_1)$ into $C(K_2)$.

Conversely, if $\varepsilon \in \{-1, 1\}$ and $\vartheta : K_2 \rightarrow K_1$ is a homeomorphism, then $f \mapsto \varepsilon(f \circ \vartheta)$ is an isometric isomorphism between $C(K_1)$ and $C(K_2)$ that maps $\mathbb{R}\mathbf{1}_1$ onto $\mathbb{R}\mathbf{1}_2$, and hence it induces an isometric isomorphism between the respective quotient spaces $\overline{C(K_1)}$ and $\overline{C(K_2)}$. □

Our next goal is to describe the group of surjective isometries from $\overline{C(K)}$ to itself. For any real normed space X , the Mazur–Ulam theorem shows that any surjective isometry from X to X is the composition of an isometric isomorphism and a translation. Clearly the subgroup of isometric isomorphisms and the subgroup of translations have trivial intersection, and it is easily verified that a translation by $x \in X$ conjugated by an isometric isomorphism T of X yields a translation by Tx . This shows that the group of surjective isometries from X to X is a semidirect product of these two subgroups. In the case of $X = \overline{C(K)}$, we know the group of isometric isomorphisms by Theorem 4.5 which yields the following description.

Proposition 4.6. *If K be a compact Hausdorff space, then the group of surjective isometries of $\overline{C(K)}$ is isomorphic to $\overline{C(K)} \rtimes (C_2 \times \text{Homeo}(K))$ if and only if $|K| \geq 3$, where C_2 is the cyclic group of order 2.*

Note that in the above proposition, $\vartheta \in \text{Homeo}(K)$ acts on $\overline{f} \in \overline{C(K)}$ as $\overline{f \circ \vartheta^{-1}}$ (not as $\overline{f \circ \vartheta}$).

Proof. If K has only two elements, then multiplication by -1 coincides with the non-trivial homeomorphism, and if $|K| \leq 1$, then multiplication by -1 is the identity. In all other cases there is no overlap, and the result follows from the above discussion. □

Translating Theorem 4.5 and Proposition 4.6 back to $(P(C(K_i)_+^\circ), d_H)$ through the pointwise exponential (the inverse of the pointwise logarithm) yields the following.

Theorem 4.7. *Let K_1, K_2 be compact Hausdorff spaces. A map*

$$h : (P(C(K_1)_+^\circ), d_H) \rightarrow (P(C(K_2)_+^\circ), d_H)$$

is a surjective isometry if and only if there exist $g \in C(K_2)_+^\circ$, $\varepsilon \in \{-1, 1\}$, and a homeomorphism $\vartheta : K_2 \rightarrow K_1$ such that

$$h(\bar{f}) = \overline{g \cdot (f \circ \vartheta)^\varepsilon} \quad \text{for all } \bar{f} \in P(C(K_2)_+^\circ).$$

If $K_1 = K_2 = K$ and $|K| \geq 3$, then the isometry group is given by

$$\text{Isom}(P(C(K)_+^\circ), d_H) \cong \overline{C(K)} \rtimes (C_2 \times \text{Homeo}(K)),$$

where C_2 is the cyclic group of order 2.

If μ_i ($i \in \{1, 2\}$) is a strictly positive measure on K_i , and we identify $P(C(K_i)_+^\circ)$ with $\Delta(K_i, \mu_i)$, we obtain Theorem 1.2.

Remark 4.8. In the above theorem, everything is projectively linear except for the inversion. It follows that if K is a compact Hausdorff space with at least three elements, the index of the collineation group in the isometry group $\text{Isom}(P(C(K)_+^\circ), d_H)$ equals 2.

Theorem 4.7 has the following interesting consequence.

Corollary 4.9. *If K_1, K_2 are compact Hausdorff spaces, then $(P(C(K_1)_+^\circ), d_H)$ and $(P(C(K_2)_+^\circ), d_H)$ are isometric if and only if K_1 and K_2 are homeomorphic.*

It would be interesting to study non-commutative versions of Theorem 4.7. In particular, one could look at Hilbert’s metric isometries on the interior of the cone A_+ of positive self-adjoint elements in a unital C^* -algebra. In view of the characterisation of Thompson’s metric isometries on A_+° by Hatori and Molnár [9], it seems plausible that each Hilbert’s metric isometry $h : P(A_+^\circ) \rightarrow P(A_+^\circ)$ is of the form $h(\bar{a}) = \overline{h(e)^{1/2} J(a^\varepsilon) h(e)^{1/2}}$, where $\varepsilon \in \{-1, 1\}$ is fixed, e is the unit in A , and J is a Jordan* isomorphism.

More generally it seems worthwhile to investigate if Walsh’s results in [24] can be extended to infinite dimensions. In [24] Walsh showed that every isometry of a finite dimensional Hilbert geometry is a projective linear automorphism, except when the domain comes from a non-Lorentzian symmetric cone. It is well known that the symmetric cones in finite dimensional vector spaces are precisely the interiors of the cones of squares of Euclidean Jordan algebras by the fundamental work of Koecher [11] and Vinberg [25]. Thus, Walsh’s result provides a link between the existence of an isometry of (Ω, d_H) that is not a projective linear automorphism and a Jordan algebra structure on the vector space above Ω . It might well be true

that in a general order unit space (X, C, u) we have that the existence of a Hilbert's metric isometry on $P(C^\circ)$ that is not a projective linear automorphism implies that X has a Jordan algebra structure and C is the cone of squares.

Acknowledgment

B.L. was supported by EPSRC grant EP/J008508/1. M.R. was supported by EPSRC grant EP/J500446/1 and M.W. was supported by EPSRC grant EP/J008508/1.

References

1. C. D. Aliprantis and R. Tourky, *Cones and Duality*. Graduate Studies in Mathematics, Vol. 84 (Amer. Math. Soc., 2007).
2. G. Birkhoff, Extensions of Jentzsch's theorems. *Trans. Amer. Math. Soc.* **85** (1957) 219–277.
3. A. Bosché, Symmetric cones, the Hilbert and Thompson metrics, arXiv:1207.3214.
4. M. Burger, A. Iozzi and N. Monod, Equivariant embeddings of trees into hyperbolic spaces, *Int. Math. Res. Not.* **22** (2005) 1331–1369.
5. H. Busemann and P. J. Kelly, *Projective Geometry and Projective Metrics* (Academic Press, 1953).
6. J. B. Conway, *A Course in Functional Analysis*, 2nd edn., Graduate Texts in Mathematics, Vol. 96 (Springer-Verlag, 1990).
7. P. de la Harpe, On Hilbert's metric for simplices, in *Geometric Group Theory*, Vol. 1, Sussex, 1991, London Math. Soc. Lecture Note Ser., Vol. 181 (Cambridge Univ. Press, 1993), pp. 97–119.
8. S. Francaviglia and A. Martino, The isometry group of outer space, *Adv. Math.* **231** (2012) 1940–1973.
9. O. Hatori and L. Molnár, Isometries of the unitary groups and Thompson isometries of the spaces of invertible positive elements in C^* -algebras. *J. Math. Anal. Appl.* **409** (2014) 158–167.
10. D. Hilbert, Über die gerade Linie als kürzeste Verbindung zweier Punkte, *Math. Ann.* **46** (1895) 91–96.
11. M. Koecher, Positivitätsbereiche im \mathbb{R}^n , *Amer. J. Math.* **79** (1957) 575–596.
12. B. Lemmens and R. Nussbaum, *Nonlinear Perron–Frobenius Theory*, Cambridge Tracts in Mathematics, Vol. 189 (Cambridge Univ. Press, 2012).
13. B. Lemmens and R. Nussbaum, Birkhoff's version of Hilbert's metric and applications, in *Handbook of Hilbert Geometry*, eds. A. Papadopoulos and M. Troyanov, IRMA Lectures in Mathematics and Theoretical Physics, Vol. 22 (European Math. Soc., 2014), pp. 275–306.
14. B. Lemmens and M. Roelands, Unique geodesics for Thompson's metric, *Ann. Inst. Fourier (Grenoble)* **65** (2015) 315–348.
15. B. Lemmens and C. Walsh, Isometries of polyhedral Hilbert geometries, *J. Topol. Anal.* **3** (2011) 213–241.
16. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces, II. Function Spaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Vol. 97 (Springer-Verlag, 1979).
17. V. Matveev and M. Troyanov, Isometries of two dimensional Hilbert metrics, *Enseign. Math.* to appear, arXiv:1409.5611.
18. L. Molnár, Thompson isometries of the space of invertible positive operators, *Proc. Amer. Math. Soc.* **137** (2009) 3849–3859.

19. N. Monod and P. Py, An exotic deformation of the hyperbolic space, *Amer. J. Math.* **136** (2014) 1249–1299.
20. R. D. Nussbaum, Hilbert’s projective metric and iterated nonlinear maps, *Mem. Amer. Math. Soc.* **75** (1988).
21. A. Papadopoulos and M. Troyanov, *Handbook of Hilbert Geometry*, IRMA Lectures in Mathematics and Theoretical Physics, Vol. 22 (European Math. Soc., 2014).
22. T. Speer, Isometries of the Hilbert metric, arXiv:1411.1826.
23. A. C. Thompson, On certain contraction mappings in a partially ordered vector space, *Proc. Amer. Math. Soc.* **14** (1963) 438–443.
24. C. Walsh, Gauge-reversing maps on cones, and Hilbert and Thompson isometries, arXiv:1312.7871.
25. E. B. Vinberg, Homogeneous cones, *Sov. Math. Dokl.* **1** (1960) 787–790.